

# Logistic Regression

Learning Centre

# Content

---

1

What is logistic  
regression?

2

Types of  
logistic  
regression

3

Worked  
example on  
binary logistic  
regression

# What is Logistic Regression?

---

Like the multiple regression, logistic regression is a statistical analysis used to examine relationships between independent variables (predictors) and a dependant variable (criterion)

The main difference is in logistic regression, the criterion is *nominal* (predicting group membership). For example, do age and gender predict whether one signs up for swimming lessons (yes/no)

# Types of Logistic Regression

---

- There are primarily 2 types of logistic regression: (1) Binary and (2) Multinomial models. The difference lies in the types of the criterion variable
- Binary logistic regression is for a dichotomous criterion (i.e., 2-level variable)
- Multinomial logistic regression is for a multicategorical criterion (i.e., a variable with more than 2 levels)
- This set of slides focuses on binary logistic regression

# Example...

A researcher would like to find out if the three predictors can predict successful enrolment into the Masters of Psychology program at JCU. The researcher recruited 30 participants who applied for the program, and asked them the following questions:

1. Interest in the Masters of Psychology program (rated 1-100)
2. Average overall score from a previous degree (scored 1-100)
3. Holding a psychology degree (yes/no)
4. Successful enrolment (yes/no)

A binary logistic regression was then conducted.

*Note that sample size of 30 was used only for illustration purposes, an actual study would require larger sample size!*

# Location of SPSS Data Files for Practice

---



Example SPSS data for practice are available on LearnJCU:

Log in to LearnJCU -> Organisations -> Learning Centre JCU Singapore ->  
Statistics Support -> Statistics Resources -> SPSS Data for Practice

# Assumptions Testing

---

01

Multicollinearity

03

**Logit Linearity**

02


**Independence of  
errors**

04

Outliers

# Assumptions Testing

Please refer to the SPSS guide on **Multiple Regression** of how to conduct the four assumption tests at <https://www.jcu.edu.sg/current-students/student-support-services/learning-support/statistics-and-mathematics-support>



- ANOVAs
- Chi-Square Tests
- Correlation
- MANOVA
- **Multiple Regression**
- t-tests



# Assumptions Testing

---

01

Multicollinearity

03

**Logit Linearity**

02

**Independence of  
errors**

04

Outliers


# Assumptions Testing: Logit Linearity

This is an assumption that the relationship between each continuous predictor and a criterion is linear.

- *Interest* and *PreviousScore* are continuous, thus they have to be tested for this assumption.
- *PsychDegree* is categorical, hence it is not required to be tested.


To test for this, we first need to create new variables in our dataset: Logit functions of the continuous IVs

- Transform → Compute Variable




# Assumptions Testing: Logit Linearity

1. Select 'Arithmetic' under Function group, and double click on 'Ln' under Functions and special variables
2. LN should appear under Numeric Expression
3. To create the logit expression of the first continuous variable (*Interest*), double click on Interest
4. Name the target variable – LnInterest
5. OK



# Assumptions Testing: Logit Linearity

Repeat the procedures, this time creating the logit function of the other continuous variable (*PreviousScore*)




# Assumptions Testing: Logit Linearity

You will see these 2 new variables in your dataset

Serialnum ber	Interest	PreviousS core	PsychDe gree	Enrolled	LnInterest	LnPreviousScore
1	73	50	1	1	4.29	3.91
2	89	50	1	1	4.49	3.91
3	89	50	1	1	4.49	3.91
4	94	50	1	1	4.54	3.91
5	77	50	1	1	4.34	3.91
6	65	60	1	2	4.17	4.09
7	69	60	1	2	4.23	4.09
8	55	55	1	2	4.01	4.01
9	81	60	1	2	4.39	4.09
10	75	70	1	1	4.32	4.25
11	69	70	1	1	4.23	4.25
12	70	65	1	1	4.25	4.17
13	93	68	1	1	4.53	4.22
14	79	69	1	1	4.37	4.23
15	70	70	1	1	4.25	4.25
16	90	89	1	2	4.50	4.49
17	73	75	1	2	4.29	4.32
18	80	80	1	2	4.38	4.38
19	86	79	1	2	4.45	4.37
20	78	78	1	2	4.36	4.36
21	82	77	2	2	4.41	4.34
22	81	68	2	1	4.39	4.22
23	78	70	2	1	4.36	4.25
24	76	71	2	1	4.33	4.26
25	96	80	2	2	4.56	4.38
26	72	68	2	1	4.28	4.22
27	65	75	2	2	4.17	4.32
28	66	77	2	2	4.19	4.34
29	75	80	2	2	4.32	4.38
30	70	82	2	2	4.25	4.41


# Assumptions Testing: Logit Linearity

To conduct the assumption test for logit linearity, go to **Analyze -> Regression -> Binary Logistic**




# Assumptions Testing: Logit Linearity

1. Move 'Enrolled' into the Dependent box
2. Move 'Interest' and 'PreviousScore' into the Covariates box
3. Holding the Ctrl key, then select 'Interest' and 'LnInterest', and click on '>a\*b>' to enter the interaction term into the Covariates box
4. Repeat Step 3 for 'PreviousScore' and 'LnPreviousScore'



# Assumptions Testing: Logit Linearity

- You should have 4 Covariates in total
- Click OK





# Assumptions Testing: Logit Linearity

Variables in the Equation		B	S.E.	Wald	df	Sig.	Exp(B)
Step 1 <sup>a</sup>	Interest	-8.430	6.888	1.498	1	.221	.000
	PreviousScore	-8.936	5.091	3.081	1	.079	.000
	Interest by LnInterest	1.550	1.282	1.463	1	.227	4.712
	LnPreviousScore by PreviousScore	1.761	.989	3.172	1	.075	5.821
	Constant	230.910	129.181	3.195	1	.074	1.918E+100


a. Variable(s) entered on step 1: Interest, PreviousScore, Interest \* LnInterest, LnPreviousScore \* PreviousScore .

Since the  $p$  values of the interaction terms are above .05, we conclude that the assumption for logit linearity is *not* violated

# Logistic Regression

Now to conduct the main analysis...


Analyze -> Regression ->  
Binary Logistic



# Logistic Regression


1. Move 'Enrolled' into the Dependent box
2. Move 'Interest', 'PreviousScore' and 'PsychDegree' into the Covariates box

*Note that 'PsychDegree' is a categorical variable.*




# Logistic Regression

3. Click on *Categorical*
4. Select 'PsychDegree' as a categorical covariate
5. Continue



# Logistic Regression

6. Click on **Save**
7. Select *Probabilities*, *Group membership*, *Cook's* (this can be used to screen for outliers), and *Standardized Residuals*
8. Continue



The image shows the 'Logistic Regression: Save' dialog box in SPSS. It has a title bar with the SPSS icon and the text 'Logistic Regression: Save' and a close button. The dialog is divided into several sections: 'Predicted Values' with checkboxes for 'Probabilities' and 'Group membership' (both checked); 'Influence' with checkboxes for 'Cook's' (checked), 'Leverage values', and 'DfBeta(s)'; 'Residuals' with checkboxes for 'Unstandardized', 'Logit', 'Studentized', 'Standardized' (checked), and 'Deviance'; and 'Export model information to XML file' with a text box and a 'Browse' button. At the bottom, there is a checkbox for 'Include the covariance matrix' (checked) and three buttons: 'Continue', 'Cancel', and 'Help'.

Logistic Regression: Save

**Predicted Values**

- ☒ Probabilities
- ☒ Group membership

**Influence**

- ☒ Cook's
- ☐ Leverage values
- ☐ DfBeta(s)

**Residuals**

- ☐ Unstandardized
- ☐ Logit
- ☐ Studentized
- ☒ Standardized
- ☐ Deviance

**Export model information to XML file**


**Browse**

☒ Include the covariance matrix

**Continue** **Cancel** **Help**

# Logistic Regression

9. Click on **Options**
10. Select *Classification plots*, *Hosmer-Lemeshow goodness-of-fit*, and *CI for  $\exp(B)$*
11. Continue, and OK



The image shows the 'Logistic Regression: Options' dialog box in SPSS. The 'Statistics and Plots' section is expanded, showing the following options:

- ☒ Classification plots
- ☒ Hosmer-Lemeshow goodness-of-fit
- ☐ Casewise listing of residuals
- ☐ Correlations of estimates
- ☐ Iteration history
- ☒ CI for  $\exp(B)$  (95 %)
- ☒ Outliers outside 2 std. dev.
- ☐ All cases

The 'Display' section shows:

- ☒ At each step
- ☐ At last step

The 'Probability for Stepwise' section shows:

- Entry: 0.05
- Removal: 0.10

The 'Classification cutoff' is set to 0.5, and 'Maximum iterations' is set to 20.

Other options include:

- ☐ Conserve memory for complex analyses or large datasets
- ☒ Include constant in model

Buttons at the bottom: Continue, Cancel, Help.

# Assumptions Testing: Outliers

---

01

Multicollinearity

03

**Logit Linearity**

02

**Independence of  
errors**

04

Outliers

# Assumptions Testing: Outliers

- Outliers can be tested together with the main analysis
- Looking at the dataset, *Cook's distance* is added as a new variable
- Since all the values are  $< 1$ , we conclude that there are no outliers

Serialnum ber	Interest	PreviousS core	PsychDe gree	Enrolled	LnInterest	LnPreviousScore	PRE_1	PGR_1	COO_1	ZRE_1
1	73	50	1	1	4.29	3.91	.07239	1	.00892	-.27935
2	89	50	1	1	4.49	3.91	.02224	1	.00157	-.15083
3	89	50	1	1	4.49	3.91	.02224	1	.00157	-.15083
4	94	50	1	1	4.54	3.91	.01524	1	.00092	-.12441
5	77	50	1	1	4.34	3.91	.05423	1	.00565	-.23946
6	65	60	1	2	4.17	4.09	.41803	1	.23694	1.17990
7	69	60	1	2	4.23	4.09	.34547	1	.24950	1.37645
8	55	55	1	2	4.01	4.01	.41044	1	.68618	1.19849
9	81	60	1	2	4.39	4.09	.17314	1	.63564	2.18530
10	75	70	1	1	4.32	4.25	.62296	2	.15271	-1.28539
11	69	70	1	1	4.23	4.25	.72399	2	.28544	-1.61960
12	70	65	1	1	4.25	4.17	.52140	2	.11212	-1.04375
13	93	68	1	1	4.53	4.22	.23053	1	.08954	-.54736
14	79	69	1	1	4.37	4.23	.50840	2	.10924	-1.01695
15	70	70	1	1	4.25	4.25	.70834	2	.25187	-1.55840
16	90	89	1	2	4.50	4.49	.91629	2	.01731	.30226
17	73	75	1	2	4.29	4.32	.81121	2	.02494	.48242
18	80	80	1	2	4.38	4.38	.84817	2	.02401	.42310
19	86	79	1	2	4.45	4.37	.74983	2	.06942	.57761
20	78	78	1	2	4.36	4.36	.82545	2	.02594	.45985
21	82	77	2	2	4.41	4.34	.60281	2	.10282	.81172
22	81	68	2	1	4.39	4.22	.27912	1	.07000	-.62226
23	78	70	2	1	4.36	4.25	.40203	1	.10982	-.81996
24	76	71	2	1	4.33	4.26	.47936	1	.14367	-.95955
25	96	80	2	2	4.56	4.38	.45503	1	.57961	1.09438
26	72	68	2	1	4.28	4.22	.43649	1	.14852	-.88010
27	65	75	2	2	4.17	4.32	.80317	2	.04636	.49504
28	66	77	2	2	4.19	4.34	.83887	2	.03130	.43827
29	75	80	2	2	4.32	4.38	.80807	2	.03018	.48736
30	70	82	2	2	4.25	4.41	.89505	2	.01390	.34243



# Logistic Regression: Results

Classification Table<sup>a,b</sup>

			Predicted		Percentage Correct
			Enrolled No	Enrolled Yes	
Step 0	Enrolled	No	0	15	.0
		Yes	0	15	100.0
	Overall Percentage				50.0

a. Constant is included in the model.

b. The cut value is .500

- The purpose of logistic regression is thus to find out if the *prediction accuracy* of the model can be improved by predictor variables
- This table shows the regression model with no predictors involved (block 0). This model (at Step 0) can correctly predict if someone successfully enrolled 50% of the time.

# Logistic Regression: Results

## Block 1: Method = Enter

### Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	11.598	3	.009
	Block	11.598	3	.009
	Model	11.598	3	.009

### Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	29.991 <sup>a</sup>	.321	.428

a. Estimation terminated at iteration number 5 because parameter estimates changed by less than .001.

### Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	11.809	8	.160

In block 1, all the predictors were entered simultaneously

A  $p$  value  $< .05$  suggests that, overall, the predictors significantly improved the prediction accuracy of the model

$R$  square values of the regression model

Measure of model fit. A  $p$  value  $> .05$  suggests a good model fit

# Logistic Regression: Results

Classification Table<sup>a</sup>

			Predicted		Percentage Correct
			Enrolled 0	No	
Step 1	Enrolled	0	10	5	66.7
		No	5	10	66.7
	Overall Percentage				66.7

a. The cut value is .500

In Step 1, the addition of the predictors resulted in the model being able to predict successful enrolment 66.7% of the time (compared to 50% in block 0; 16.7% improvement !)

# Logistic Regression: Results

Variables in the Equation								
		B	S.E.	Wald	df	Sig.	Exp(B)	95% C.I. for EXP(B) Lower Upper
Step 1 <sup>a</sup>	Interest	-.077	.054	2.051	1	.152	.926	.833 1.029
	PreviousScore	.160	.066	5.972	1	.015	1.174	1.032 1.335
	PsychDegree(1)	.668	1.003	.443	1	.506	1.950	.273 13.938
	Constant	-5.611	4.931	1.295	1	.255	.004	

a. Variable(s) entered on step 1: Interest, PreviousScore, PsychDegree.

- This table tells us which predictors are significant. Only Previous score is a significant predictor ( $p < .05$ )
- In logistic regression, Exp(B) is commonly used to interpret results, and is expressed as **an odds ratio**
- In other words, an increase of 1 unit in Previous score results in a 17.4% more chance of enrolling in the masters program ( $1.174 - 1 = .174$ , meaning .174 above 1)
- The other statistics (e.g., B, Wald, 95% CI) can also be reported in the writeup

# Write-Up

---

An example write-up can be found on page 228 in

**Allen, P., Bennett, K., & Heritage, B. (2019). *SPSS Statistics: A Practical Guide* (4th ed.). Cengage Learning.**

# Questions?

[learningcentre-singapore@jcu.edu.au](mailto:learningcentre-singapore@jcu.edu.au)